

Explicit product ensembles for separable quantum states

Rüdiger Schack^{*(a,b)} and Carlton M. Caves^{†(a)}

^(a)Center for Advanced Studies, Department of Physics and Astronomy,
University of New Mexico, Albuquerque, NM 87131–1156, USA

^(b)Department of Mathematics, Royal Holloway,
University of London, Egham, Surrey TW20 0EX, UK

February 1, 2008

Abstract

We present a general method for constructing pure-product-state representations for density operators of N quantum bits. If such a representation has nonnegative expansion coefficients, it provides an explicit separable ensemble for the density operator. We derive the condition for separability of a mixture of the Greenberger-Horne-Zeilinger state with the maximally mixed state.

1 Introduction

The state of a quantum system composed of N subsystems is *separable* if it can be written as a classical mixture or classical *ensemble* of tensor product states. Although it is straightforward to decide whether a pure state is separable, the same question is in general unsolved for mixed states. A simple separability criterion is known only for two subsystems, and even then only if one of the subsystems is a two-level system (qubit) and the other one is at most a three-level system (qutrit) [1, 2]

Recently it has been shown that there exists a neighborhood of the maximally mixed state in which all states are separable [3]. For N qubits the size of this neighborhood decreases exponentially with the number of qubits [4]. These results are interesting, among other reasons, because NMR quantum computers [5, 6, 7] operate with states near the maximally mixed state.

In this paper we develop a general method for constructing pure-product-state representations for N -qubit density operators; when the expansion coefficients in such a representation are nonnegative, it provides an explicit product ensemble for the density operator,

*E-mail: r.schack@rhbnc.ac.uk

†E-mail: caves@tangelo.phys.unm.edu

and the state is separable. In Sec. 2.1, we introduce a superoperator formalism that allows us to treat different discrete and continuous representations on the same footing. In Sec. 2.2, we apply the formalism to a discrete overcomplete operator basis, leading to a simple rederivation of the separability bound found in [4]. Section 2.3 introduces a “canonical” continuous representation for an arbitrary density operator, and Sec. 2.4 develops a method for deriving discrete representations from the continuous representation. In Sec. 3, we show that the canonical continuous representation is the unique representation containing only scalar and vector spherical-harmonic components; all other representations can be obtained by adding arbitrary higher-order spherical harmonics. Section 3.2 applies the canonical continuous representation to mixtures of the N -qubit “cat state” with the maximally mixed state. In Sec. 4, we derive the conditions for separability of a mixture of the Greenberger-Horne-Zeilinger state [8] with the maximally mixed state. Section 5 concludes with a brief discussion.

2 Product-state representations and separable ensembles

2.1 Superoperator formalism

The set of linear operators acting on a D -dimensional Hilbert space $\mathcal{H}$ is a D^2 -dimensional complex vector space $\mathcal{L}(\mathcal{H})$. Let us introduce operator “kets” $|A\rangle = A$ and “bras” $\langle A| = A^\dagger$, distinguished from vector kets and bras by the use of round brackets [9]. Then the natural inner product on $\mathcal{L}(\mathcal{H})$, the trace-norm inner product, can be written as $\langle A|B\rangle = \text{tr}(A^\dagger B)$. The notation $\mathcal{S} = |A\rangle\langle B|$ defines a superoperator $\mathcal{S}$ acting like

$$\mathcal{S}|X\rangle = |A\rangle\langle B|X\rangle = \text{tr}(B^\dagger X)A. \quad (1)$$

Now let the set $\{|N_j\rangle\}$ constitute a (complete or overcomplete) operator basis; i.e., let the operator kets $|N_j\rangle$ span the vector space $\mathcal{L}(\mathcal{H})$. It follows that the superoperator $\mathcal{G}$ defined by

$$\mathcal{G} \equiv \sum_j |N_j\rangle\langle N_j| \quad (2)$$

is invertible. The operators

$$Q_j \equiv \mathcal{G}^{-1}|N_j\rangle \quad (3)$$

form a dual basis, which gives rise to the following resolutions of the superoperator identity:

$$\mathbf{1} = \sum_j |Q_j\rangle\langle N_j| = \sum_j |N_j\rangle\langle Q_j|. \quad (4)$$

An arbitrary operator A can be expanded as

$$A = \sum_j |N_j\rangle\langle Q_j|A\rangle = \sum_j N_j \text{tr}(Q_j^\dagger A) \quad (5)$$

and

$$A = \sum_j |Q_j\rangle\langle N_j|A\rangle = \sum_j Q_j \text{tr}(N_j^\dagger A). \quad (6)$$

These expansions are unique if and only if the operators N_j are linearly independent.

2.2 A discrete representation for the 6 cardinal directions

The Hermitian operators $P_{j\mu}$, $j = 1, 2, 3$, $\mu = 0, 1$, defined by

$$P_{j\mu} \equiv \frac{1}{2} \left(1 + (-1)^\mu \sigma_j \right), \quad (7)$$

are the pure-state projectors corresponding to the 6 cardinal directions on the Bloch sphere. They form an overcomplete basis in the space of operators acting on a qubit. The super-operator

$$\mathcal{G} = \sum_{j,\mu} |P_{j\mu}\rangle \langle P_{j\mu}| = \frac{1}{2} \left(3|1\rangle \langle 1| + \sum_j |\sigma_j\rangle \langle \sigma_j| \right) \quad (8)$$

has normalized eigenoperators $|1\rangle/\sqrt{2}$ and $|\sigma_j\rangle/\sqrt{2}$, with eigenvalues 3 and 1, respectively, from which it follows that

$$\mathcal{G}^{-1} = \frac{1}{2} \left(\frac{1}{3} |1\rangle \langle 1| + \sum_j |\sigma_j\rangle \langle \sigma_j| \right). \quad (9)$$

The last equation allows us to find the dual-basis operators,

$$Q_{j\mu} = \mathcal{G}^{-1} |P_{j\mu}\rangle = \frac{1}{6} \left(1 + 3(-1)^\mu \sigma_j \right). \quad (10)$$

Any density operator $\rho = \frac{1}{2}(1 + \vec{S} \cdot \vec{\sigma})$ for one qubit can be represented in the form

$$\rho = \sum_{j,\mu} |P_{j\mu}\rangle \langle Q_{j\mu}| \rho = \frac{1}{6} \sum_{j,\mu} P_{j\mu} \left(1 + 3(-1)^\mu S_j \right). \quad (11)$$

For an N -qubit density operator, the analogous representation is

$$\rho = \sum_{j_1, \mu_1, \dots, j_N, \mu_N} w^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N) P_{j_1 \mu_1} \otimes \dots \otimes P_{j_N \mu_N}, \quad (12)$$

where

$$\begin{aligned} w^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N) &\equiv \text{tr}(\rho Q_{j_1 \mu_1} \otimes \dots \otimes Q_{j_N \mu_N}) \\ &= \frac{1}{6^N} \text{tr} \left(\rho \left(1 + 3(-1)^{\mu_1} \sigma_{j_1} \right) \otimes \dots \otimes \left(1 + 3(-1)^{\mu_N} \sigma_{j_N} \right) \right). \end{aligned} \quad (13)$$

This representation exists for arbitrary density operators, entangled or separable. If the coefficients $w^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N)$ are nonnegative, the density operator ρ is separable, and the representation (12) provides an explicit product-state ensemble for the density operator. The expansions (11) and (12) are not unique because the 6 operators $P_{j\mu}$ are not linearly independent. Throughout the paper we use the superscript “can,” standing for *canonical*, to denote the natural, but generally not unique expansion coefficients that come from using the procedure of Sec. 2.1.

For any N -qubit density operator ρ , the coefficients (13) obey the bound

$$w^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N) \geq \left(\text{smallest eigenvalue of } Q_{j_1 \mu_1} \otimes \dots \otimes Q_{j_N \mu_N} \right) = -\frac{2^{2N-1}}{6^N}. \quad (14)$$

This follows from the fact that $Q_{j\mu}$ has eigenvalues $2/3$ and $-1/3$. The most negative eigenvalue of the product operator $Q_{j_1 \mu_1} \otimes \dots \otimes Q_{j_N \mu_N}$ is therefore $(-1/3)(2/3)^{N-1} = -2^{2N-1}/6^N$, from which Eq. (14) follows.

Consider now an N -qubit density operator that is a mixture of the maximally mixed density operator, $1/2^N$, and an arbitrary density operator ρ_1 :

$$\rho_\epsilon = \frac{1-\epsilon}{2^N} 1 + \epsilon \rho_1. \quad (15)$$

Any density operator can be written in this form [4]. The coefficients (13) for ρ_ϵ are given by

$$\begin{aligned} w_\epsilon^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N) &= \frac{1-\epsilon}{6^N} + \epsilon w_1^{\text{can}}(j_1, \mu_1, \dots, j_N, \mu_N) \\ &\geq \frac{1-\epsilon(1+2^{2N-1})}{6^N}, \end{aligned} \quad (16)$$

which is nonnegative for

$$\epsilon \leq \frac{1}{1+2^{2N-1}}. \quad (17)$$

This bound, derived in [4] using a different route, implies that, for $\epsilon \leq 1/(1+2^{2N-1})$, all density operators of the form (15) are separable. It sets a lower bound on the size of the separable neighborhood surrounding the maximally mixed state.

2.3 Canonical continuous representation

The set of all pure-state projectors on the Bloch sphere,

$$P_{\vec{n}} \equiv |\vec{n}\rangle\langle\vec{n}| = \frac{1}{2}(1 + \vec{\sigma} \cdot \vec{n}), \quad (18)$$

forms an overcomplete basis. The corresponding superoperator

$$\mathcal{G} = \int d\Omega |P_{\vec{n}}\rangle\langle P_{\vec{n}}| = \pi \left(|1\rangle\langle 1| + \frac{1}{3} \sum_j |\sigma_j\rangle\langle\sigma_j| \right) \quad (19)$$

is proportional to the superoperator of Eq. (8). The normalized eigenoperators of $\mathcal{G}$ are $|1\rangle/\sqrt{2}$ and $|\sigma_j\rangle/\sqrt{2}$, with eigenvalues 2π and $2\pi/3$, respectively. Thus $\mathcal{G}^{-1}$ can be expressed as

$$\mathcal{G}^{-1} = \frac{1}{4\pi} \left(|1\rangle\langle 1| + 3 \sum_j |\sigma_j\rangle\langle\sigma_j| \right). \quad (20)$$

The dual-basis operators are then given by

$$Q_{\vec{n}} = \mathcal{G}^{-1}|P_{\vec{n}}) = \frac{1}{4\pi}(1 + 3\vec{\sigma} \cdot \vec{n}) . \quad (21)$$

Any density operator $\rho = \frac{1}{2}(1 + \vec{S} \cdot \vec{\sigma})$ for one qubit can be represented in the form

$$\rho = \int d\Omega |P_{\vec{n}})(Q_{\vec{n}}|\rho) = \frac{1}{4\pi} \int d\Omega P_{\vec{n}}(1 + 3\vec{S} \cdot \vec{n}) . \quad (22)$$

For N qubits, we define the pure-product-state projector

$$P(\tilde{n}) \equiv P_{\vec{n}_1} \otimes \cdots \otimes P_{\vec{n}_N} = \frac{1}{2^N}(1 + \vec{\sigma} \cdot \vec{n}_1) \otimes \cdots \otimes (1 + \vec{\sigma} \cdot \vec{n}_N) , \quad (23)$$

where $\tilde{n}$ stands for the collection of unit vectors $\vec{n}_1, \dots, \vec{n}_N$. Any N -qubit density operator can be expanded as

$$\rho = \int d\Omega_{\tilde{n}} w^{\text{can}}(\tilde{n}) P(\tilde{n}) \equiv \int d\Omega_{n_1} \cdots d\Omega_{n_N} w^{\text{can}}(\tilde{n}) P(\tilde{n}) , \quad (24)$$

where

$$w^{\text{can}}(\tilde{n}) \equiv \text{tr}(\rho Q(\tilde{n})) \quad (25)$$

is the canonical expansion function, with

$$Q(\tilde{n}) \equiv Q_{\vec{n}_1} \otimes \cdots \otimes Q_{\vec{n}_N} = \frac{1}{(4\pi)^N}(1 + 3\vec{\sigma} \cdot \vec{n}_1) \otimes \cdots \otimes (1 + 3\vec{\sigma} \cdot \vec{n}_N) . \quad (26)$$

If the coefficients $w^{\text{can}}(\tilde{n})$ are nonnegative, the density operator ρ is separable, and the representation (24) provides an explicit product-state ensemble for the density operator.

Using the same argument as in the preceding subsection, one can show that for any density operator ρ [4],

$$w^{\text{can}}(\tilde{n}) \geq \left(\begin{array}{c} \text{smallest eigenvalue} \\ \text{of } Q(\tilde{n}) \end{array} \right) = -\frac{2^{2N-1}}{(4\pi)^N} . \quad (27)$$

Applying this result to density operators of the form (15), one arrives again at the lower bound (17) on the size of the separable neighborhood of the maximally mixed density operator [4].

2.4 Other discrete representations

The expansion coefficients (13) for the discrete representation of Sec. 2.2 can be obtained by evaluating the expansion coefficients (25) for the canonical continuous representation along the cardinal directions for each qubit and then renormalizing the resulting function. This is no accident. It is easy to characterize a class of discrete representations whose expansion coefficients are similarly related to the continuous representation.

Consider a discrete set of projection operators,

$$P_{\vec{n}_\alpha} = |\vec{n}_\alpha\rangle\langle\vec{n}_\alpha| = \frac{1}{2}(1 + \vec{\sigma} \cdot \vec{n}_\alpha), \quad \alpha = 1, \dots, K. \quad (28)$$

The corresponding superoperator

$$\begin{aligned} \mathcal{G} &= \sum_{\alpha=1}^K |P_{\vec{n}_\alpha}\rangle(P_{\vec{n}_\alpha}| \\ &= \frac{1}{4} \left(K|1\rangle(1| + \sum_{\alpha} (\vec{n}_\alpha \cdot |\vec{\sigma}|)(1| + |1\rangle(\vec{\sigma}| \cdot \vec{n}_\alpha) + \sum_{j,k} |\sigma_j\rangle(\sigma_k| \sum_{\alpha} (n_{\alpha})_j (n_{\alpha})_k) \right) \end{aligned} \quad (29)$$

generates dual-basis operators and expansion coefficients proportional to those for the continuous representation if and only if $\mathcal{G}$ is proportional to the superoperator (19) for the continuous representation, i.e., if and only if

$$0 = \sum_{\alpha} \vec{n}_{\alpha}, \quad (30)$$

$$\frac{1}{3} \delta_{jk} = \frac{1}{K} \sum_{\alpha} (n_{\alpha})_j (n_{\alpha})_k. \quad (31)$$

The trace of Eq. (31) is satisfied by any collection of unit vectors.

When these conditions are satisfied, the superoperator (29) simplifies to

$$\mathcal{G} = \frac{K}{4} \left(|1\rangle(1| + \frac{1}{3} \sum_j |\sigma_j\rangle(\sigma_j|) \right), \quad (32)$$

with an inverse

$$\mathcal{G}^{-1} = \frac{1}{K} \left(|1\rangle(1| + 3 \sum_j |\sigma_j\rangle(\sigma_j|) \right) \quad (33)$$

that generates dual-basis operators

$$Q_{\vec{n}_\alpha} = \mathcal{G}^{-1} |P_{\vec{n}_\alpha}\rangle = \frac{1}{K} (1 + 3\vec{\sigma} \cdot \vec{n}_\alpha). \quad (34)$$

Any density operator $\rho = \frac{1}{2}(1 + \vec{S} \cdot \vec{\sigma})$ for one qubit can be represented in the form

$$\rho = \sum_{\alpha} |P_{\alpha}\rangle(Q_{\alpha}|\rho) = \frac{1}{K} \sum_{\alpha} P_{\alpha} (1 + 3\vec{S} \cdot \vec{n}_{\alpha}). \quad (35)$$

The conditions (30) and (31) are satisfied by unit vectors that point to the vertices of any regular polyhedron inscribed within the Bloch sphere—i.e., a tetrahedron, octahedron, cube, icosahedron, or dodecahedron. The 4 projectors for a tetrahedron are linearly independent, making the corresponding tetrahedral representation (35) unique. An octahedron gives rise to the 6-cardinal-direction representation of Sec. 2.2. The regular polyhedra by no means exhaust the possibilities for representations of this sort. One can use simultaneously the vertices from any number of polyhedra. Moreover, one can produce an appropriate set

of unit vectors by the following procedure: starting with any set of unit vectors in the first octant, reflect the set successively through the three Cartesian planes to produce a set of vectors that occupies all eight octants.

For more than one qubit, the possibilities for discrete representations that follow from the continuous representation are even more diverse. The simplest possibility is to use the same set of pure-state projectors for each qubit, but this is not necessary. One can use a different set of projectors for each qubit, or more generally, one can use a “tree structure,” in which there is a different set of projectors for the n th qubit for each projector for the first $n - 1$ qubits. There are possibilities more general even than those with a tree structure, analogous to the “domino states” introduced for two qutrits in [10]. We do not dwell here on the myriad of discrete representations of this sort, merely noting that since the expansion coefficients in such representations follow from the expansion coefficients for the continuous representation, all such representations lead to the same general lower bound (17) on the size of the separable neighborhood surrounding the maximally mixed state. Both the continuous and these discrete representations can, however, give better bounds on the separability of *particular* states of the form (15), a point to which we return in Sec. 3.2.

3 Spherical-harmonic expansions

In this section we explore the relationship between product-state representations and spherical-harmonic expansions.

3.1 Spherical-harmonic expansions and the Pauli representation

We begin by noting that if $\vec{n}$ is a unit vector, then

$$\begin{aligned}\vec{\sigma} \cdot \vec{n} &= \sigma_3 \cos \theta + \sigma_1 \sin \theta \cos \phi + \sigma_2 \sin \theta \sin \phi \\ &= \sqrt{\frac{4\pi}{3}} \left(\sigma_3 Y_1^{0*} + \sigma_1 \frac{1}{\sqrt{2}} (-Y_1^{+1*} + Y_1^{-1*}) - i\sigma_2 \frac{1}{\sqrt{2}} (Y_1^{+1*} + Y_1^{-1*}) \right) \\ &= \sigma_1^0 Y_1^{0*} + \sigma_1^{+1} Y_1^{+1*} + \sigma_1^{-1} Y_1^{-1*},\end{aligned}\tag{36}$$

where

$$Y_0^0 = \frac{1}{\sqrt{4\pi}}, \quad Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos \theta, \quad Y_1^{\pm 1} = \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\phi}\tag{37}$$

are spherical harmonics and, in the last line, we define

$$\sigma_0^0 \equiv \sqrt{4\pi} \, 1, \quad \sigma_1^0 \equiv \sqrt{\frac{4\pi}{3}} \sigma_3, \quad \sigma_1^{\pm 1} \equiv \mp \sqrt{\frac{2\pi}{3}} (\sigma_1 \pm i\sigma_2).\tag{38}$$

We can then write any pure-state projector on the Bloch sphere in the form

$$P_{\vec{n}} = |\vec{n}\rangle\langle\vec{n}| = \frac{1}{2}(1 + \vec{\sigma} \cdot \vec{n}) = \frac{1}{2} \sum_{l=0}^1 \sum_{m=-l}^l \sigma_l^m Y_l^{m*}.\tag{39}$$

Any N -qubit density operator can be written in the form

$$\rho = \int d\Omega_{\tilde{n}} w(\tilde{n}) P(\tilde{n}) , \quad (40)$$

where the expansion coefficients $w(\tilde{n}) \equiv w(\vec{n}_1, \dots, \vec{n}_N)$ are not necessarily nonnegative [the expansion coefficients $w(\tilde{n})$ need not be those of the canonical representation (25)]. A density operator ρ is separable if and only if it can be written in the form (40) for some nonnegative probability density $w(\tilde{n})$. Now expand the product projector (23) in terms of spherical harmonics,

$$P(\tilde{n}) = \frac{1}{2^N} \sum_{\{l_j \leq 1, m_j\}} Y_{l_1}^{m_1*} \cdots Y_{l_N}^{m_N*} \sigma_{l_1}^{m_1} \otimes \cdots \otimes \sigma_{l_N}^{m_N} , \quad (41)$$

where

$$\sum_{\{l_j \leq 1, m_j\}} \equiv \sum_{l_1=0}^1 \sum_{m_1=-l_1}^{l_1} \cdots \sum_{l_N=0}^1 \sum_{m_N=-l_N}^{l_N} \quad (42)$$

denotes a sum that contains only scalar and vector terms ($l = 0, 1$), and also expand the expansion coefficients,

$$w(\tilde{n}) = \sum_{\{l_j, m_j\}} a_{l_1 \dots l_N}^{m_1 \dots m_N} Y_{l_1}^{m_1} \cdots Y_{l_N}^{m_N} , \quad (43)$$

where

$$\sum_{\{l_j, m_j\}} \equiv \sum_{l_1=0}^{\infty} \sum_{m_1=-l_1}^{l_1} \cdots \sum_{l_N=0}^{\infty} \sum_{m_N=-l_N}^{l_N} \quad (44)$$

denotes an unrestricted sum over spherical harmonics. Inserting Eqs. (41) and (43) into Eq. (40) and using the orthonormality of the spherical harmonics yields the (unique) Pauli-operator representation of ρ :

$$\rho = \frac{1}{2^N} \sum_{\{l_j \leq 1, m_j\}} a_{l_1 \dots l_N}^{m_1 \dots m_N} \sigma_{l_1}^{m_1} \otimes \cdots \otimes \sigma_{l_N}^{m_N} . \quad (45)$$

We see that ρ depends on and determines only the scalar and vector parts of $a_{l_1 \dots l_N}^{m_1 \dots m_N}$; the higher-order-spherical-harmonic (HOSH) content of $w(\tilde{n})$ corresponds to the freedom in constructing pure-product-state representations of ρ . It is straightforward to generate all pure-product-state representations: given the Pauli-operator representation (45) of ρ , any $w(\tilde{n})$ is given by Eq. (43), where the coefficients $a_{l_1 \dots l_N}^{m_1 \dots m_N}$ are arbitrary except for $l_1, \dots, l_N \leq 1$.

The canonical continuous representation $w^{\text{can}}(\tilde{n})$ of Eq. (25) is the unique pure-product-state representation given by the $l = 0$ and $l = 1$ components. We can see this directly in the following way. Write the Pauli representation of ρ in the form

$$\rho = \frac{1}{2^N} \sum_{\alpha_1, \dots, \alpha_N} c_{\alpha_1 \dots \alpha_N} \sigma_{\alpha_1} \otimes \cdots \otimes \sigma_{\alpha_N} , \quad (46)$$

where

$$c_{\alpha_1 \dots \alpha_N} = \text{tr}(\rho \sigma_{\alpha_1} \otimes \cdots \otimes \sigma_{\alpha_N}) = \langle \sigma_{\alpha_1} \otimes \cdots \otimes \sigma_{\alpha_N} \rangle . \quad (47)$$

Here $\sigma_0 = 1$, and the sums over Greek indices run from 0 to 3. Now we have that

$$\begin{aligned}
w^{\text{can}}(\tilde{n}) &= \frac{1}{(4\pi)^N} \langle (1 + 3\vec{\sigma} \cdot \vec{n}_1) \otimes \cdots \otimes (1 + 3\vec{\sigma} \cdot \vec{n}_N) \rangle \\
&= \left(\frac{3}{4\pi}\right)^N \sum_{\alpha_1, \dots, \alpha_N} \langle (n_1)_{\alpha_1} \sigma_{\alpha_1} \otimes \cdots \otimes (n_N)_{\alpha_N} \sigma_{\alpha_N} \rangle \\
&= \left(\frac{3}{4\pi}\right)^N \sum_{\alpha_1, \dots, \alpha_N} c_{\alpha_1 \dots \alpha_N} (n_1)_{\alpha_1} \cdots (n_N)_{\alpha_N} ,
\end{aligned} \tag{48}$$

where $(n_j)_{\alpha_j} \equiv 1/3$ if $\alpha_j = 0$ and $(n_j)_{\alpha_j}$ is a Cartesian component of $\vec{n}_j$ if $\alpha_j = 1, 2$, or 3. Thus any pure-product-state representation of a density operator ρ can be obtained by adding HOSH to the canonical continuous representation $w^{\text{can}}(\tilde{n})$. The discrete representations discussed in Sec. 2.4 are examples—though by no means the only examples—of representations with HOSH content.

The continuous canonical representation can be generated directly from the Pauli representation of ρ by making the substitutions

$$2^N \rho \longrightarrow w^{\text{can}}(\tilde{n}) , \tag{49}$$

$$1 \longrightarrow \frac{1}{4\pi} , \quad \sigma_j \longrightarrow \frac{3}{4\pi} n_j . \tag{50}$$

These rules are equivalent to $\sigma_l^m \longrightarrow Y_l^m$ and to $\sigma_\alpha \longrightarrow (3/4\pi)n_\alpha$.

3.2 The ϵ -cat state

As an example, consider the following N -qubit generalization of the Werner [11] state,

$$\rho_\epsilon = \frac{1-\epsilon}{2^N} 1 + \epsilon |\psi_{\text{cat}}\rangle \langle \psi_{\text{cat}}| , \tag{51}$$

where

$$|\psi_{\text{cat}}\rangle \equiv \frac{1}{\sqrt{2}} (|0 \dots 0\rangle + |1 \dots 1\rangle) \tag{52}$$

is the N -qubit “cat state.” We call the mixed state (51) the ϵ -cat state.

Now let $|0\rangle$ and $|1\rangle$ be the eigenstates of σ_3 , with $|0\rangle$ at the top of the Bloch sphere and $|1\rangle$ at the bottom, so that $\sigma_1 = |0\rangle\langle 1| + |1\rangle\langle 0|$, $\sigma_2 = -i|0\rangle\langle 1| + i|1\rangle\langle 0|$, and $\sigma_3 = |0\rangle\langle 0| - |1\rangle\langle 1|$. It follows that the Pauli representation of the ϵ -cat state is

$$\begin{aligned}
\rho_\epsilon &= \frac{1}{2^N} \left((1-\epsilon) 1 \right. \\
&\quad + \frac{\epsilon}{2} (1 + \sigma_3) \otimes \cdots \otimes (1 + \sigma_3) + \frac{\epsilon}{2} (1 - \sigma_3) \otimes \cdots \otimes (1 - \sigma_3) \\
&\quad \left. + \frac{\epsilon}{2} (\sigma_1 + i\sigma_2) \otimes \cdots \otimes (\sigma_1 + i\sigma_2) + \frac{\epsilon}{2} (\sigma_1 - i\sigma_2) \otimes \cdots \otimes (\sigma_1 - i\sigma_2) \right) .
\end{aligned} \tag{53}$$

Applying the rules (49) and (50) yields the continuous canonical representation of ρ_ϵ :

$$\begin{aligned}
w_\epsilon^{\text{can}}(\tilde{n}) = & \frac{1}{(4\pi)^N} \left((1 - \epsilon) \right. \\
& + \frac{\epsilon}{2} (1 + 3 \cos \theta_1) \cdots (1 + 3 \cos \theta_N) + \frac{\epsilon}{2} (1 - 3 \cos \theta_1) \cdots (1 - 3 \cos \theta_N) \\
& \left. + \epsilon 3^N \sin \theta_1 \cdots \sin \theta_N \cos(\phi_1 + \dots + \phi_N) \right). \tag{54}
\end{aligned}$$

The minimum value of $w_\epsilon^{\text{can}}(\tilde{n})$ occurs either (i) when one qubit is evaluated at the north (south) pole, and the other $N - 1$ qubits are evaluated at the south (north) pole, or (ii) when all the qubits are evaluated on the equator with $\cos(\phi_1 + \dots + \phi_N) = -1$. For $N = 2$ and $N = 4$, both the poles and the equator give the same minimum value, which shows that the ϵ -cat state is separable for $\epsilon \leq 1/9$ for $N = 2$ and for $\epsilon \leq 1/81$ for $N = 4$. For $N = 3$ and $N = 5$, the equator gives the minimum value, which shows that the ϵ -cat state is separable for $\epsilon \leq 1/27$ for $N = 3$ [4] and for $\epsilon \leq 1/243$ for $N = 5$. For $N \geq 6$, the poles give the minimum value, which shows that ρ_ϵ is separable for

$$\epsilon \leq \frac{1}{1 \pm 2^N + 2^{2N-2}}, \tag{55}$$

the upper (lower) sign applying when N is even (odd). Notice that except for $N = 2$, all these bounds are better than the general-purpose bound of Eq. (17).

These bounds on the separability of the ϵ -cat state can be improved by using tetrahedral ensembles of the sort discussed in Sec. 2.4. The tetrahedral ensembles are chosen to have a tree structure such that the vertices avoid the places where $w_\epsilon^{\text{can}}(\tilde{n})$ has its smallest values. We do not discuss these tetrahedral bounds here, preferring instead to consider in detail the separability of the ϵ -GHZ state.

4 The ϵ -GHZ state

In this section we consider the 3-qubit version of the ϵ -cat state introduced in Sec. 3.2. In the case of 3 qubits, the cat state of Eq. (52) is called the Greenberger-Horne-Zeilinger (GHZ) state [8],

$$|\psi_{\text{GHZ}}\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle); \tag{56}$$

we call the mixture ρ_ϵ of Eq. (51) the ϵ -GHZ state. We show in this section that *the ϵ -GHZ state is separable if and only if $\epsilon \leq 1/5$* . The class of ϵ -GHZ states has been considered previously in [12].

4.1 Separability of the Werner state

Before proceeding to the proof of the separability of the ϵ -GHZ state, we review a similar, but simpler proof for the Werner state [11], the 2-qubit version of the ϵ -cat state:

$$\rho_W = \frac{1-\epsilon}{4} 1 \otimes 1 + \frac{\epsilon}{2} (|00\rangle + |11\rangle)(\langle 00| + \langle 11|)$$

$$= \frac{1}{4} \left(1 \otimes 1 + \epsilon (\sigma_3 \otimes \sigma_3 + \sigma_1 \otimes \sigma_1 - \sigma_2 \otimes \sigma_2) \right). \quad (57)$$

The Werner state is separable if and only if $\epsilon \leq 1/3$ [13, 14], a result we now show in the following way.

A density operator ρ for 2 qubits, A and B , is separable if and only if it can be written as

$$\rho = \int d\Omega_{\vec{n}_A} d\Omega_{\vec{n}_B} w(\vec{n}_A, \vec{n}_B) P_{\vec{n}_A} \otimes P_{\vec{n}_B}, \quad (58)$$

with $w(\vec{n}_A, \vec{n}_B)$ being a nonnegative probability distribution. If a state is separable, the coefficients (47) in the Pauli representation (46) of ρ become classical expectations over the distribution $w(\vec{n}_A, \vec{n}_B)$, expectations that we denote by $E[\cdot]$:

$$c_{\alpha\beta} = \text{tr}(\rho \sigma_\alpha \otimes \sigma_\beta) = \int d\Omega_{\vec{n}_A} d\Omega_{\vec{n}_B} (n_A)_\alpha (n_B)_\beta w(\vec{n}_A, \vec{n}_B) \equiv E[(n_A)_\alpha (n_B)_\beta]. \quad (59)$$

For the Werner state, the coefficients $c_{\alpha\beta}$ can be read off the Pauli representation in Eq. (57):

$$E[(n_A)_j] = E[(n_B)_j] = 0, \quad (60)$$

$$E[(n_A)_j (n_B)_k] = 0, \quad \text{for } j \neq k, \quad (61)$$

$$E[(n_A)_1 (n_B)_1] = -E[(n_A)_2 (n_B)_2] = E[(n_A)_3 (n_B)_3] = \epsilon. \quad (62)$$

If $w(\vec{n}_A, \vec{n}_B)$ is a probability distribution, we have that

$$\epsilon = \left| E[(n_A)_j (n_B)_j] \right| \leq \frac{1}{2} E[(n_A)_j^2 + (n_B)_j^2]. \quad (63)$$

Adding over the three values of j gives

$$3\epsilon \leq \frac{1}{2} E[(n_A)_1^2 + (n_A)_2^2 + (n_A)_3^2 + (n_B)_1^2 + (n_B)_2^2 + (n_B)_3^2] = 1, \quad (64)$$

which shows that if $\epsilon > 1/3$, the Werner state is nonseparable.

We still have the task of designing a product ensemble that gives the Werner state for $\epsilon \leq 1/3$. Defining three density operators

$$\rho_1 \equiv \frac{1}{2} (P_{\vec{e}_3} \otimes P_{\vec{e}_3} + P_{-\vec{e}_3} \otimes P_{-\vec{e}_3}) = \frac{1}{4} (1 \otimes 1 + \sigma_3 \otimes \sigma_3), \quad (65)$$

$$\rho_2 \equiv \frac{1}{2} (P_{\vec{e}_1} \otimes P_{\vec{e}_1} + P_{-\vec{e}_1} \otimes P_{-\vec{e}_1}) = \frac{1}{4} (1 \otimes 1 + \sigma_1 \otimes \sigma_1), \quad (66)$$

$$\rho_3 \equiv \frac{1}{2} (P_{\vec{e}_2} \otimes P_{-\vec{e}_2} + P_{-\vec{e}_2} \otimes P_{\vec{e}_2}) = \frac{1}{4} (1 \otimes 1 - \sigma_2 \otimes \sigma_2), \quad (67)$$

we see that each takes care of one of the correlations in Eq. (62). An equal mixture gives the $\epsilon = 1/3$ Werner state:

$$\frac{1}{3} (\rho_1 + \rho_2 + \rho_3) = \frac{1}{4} \left(1 \otimes 1 + \frac{1}{3} (\sigma_3 \otimes \sigma_3 + \sigma_1 \otimes \sigma_1 - \sigma_2 \otimes \sigma_2) \right). \quad (68)$$

Smaller values of ϵ can be handled by adding on a product ensemble for the maximally mixed state.

4.2 Separability of the ϵ -GHZ state

Consider now the ϵ -GHZ state

$$\begin{aligned}\rho_{\epsilon\text{GHZ}} &= \frac{1-\epsilon}{8} 1 \otimes 1 \otimes 1 + \frac{\epsilon}{2} (|000\rangle + |111\rangle) (\langle 000| + \langle 111|) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 + \epsilon \left(1 \otimes \sigma_3 \otimes \sigma_3 + \sigma_3 \otimes 1 \otimes \sigma_3 + \sigma_3 \otimes \sigma_3 \otimes 1 \right. \right. \\ &\quad \left. \left. + \sigma_1 \otimes \sigma_1 \otimes \sigma_1 - \sigma_1 \otimes \sigma_2 \otimes \sigma_2 - \sigma_2 \otimes \sigma_1 \otimes \sigma_2 - \sigma_2 \otimes \sigma_2 \otimes \sigma_1 \right) \right). \end{aligned} \quad (69)$$

A density operator ρ for 3 qubits, A , B , and C , is separable if and only if it can be written as

$$\rho = \int d\Omega_{\vec{n}_A} d\Omega_{\vec{n}_B} d\Omega_{\vec{n}_C} w(\vec{n}_A, \vec{n}_B, \vec{n}_C) P_{\vec{n}_A} \otimes P_{\vec{n}_B} \otimes P_{\vec{n}_C}, \quad (70)$$

with $w(\vec{n}_A, \vec{n}_B, \vec{n}_C)$ being a nonnegative probability distribution. If a state is separable, the coefficients (47) in the Pauli representation (46) of ρ become classical expectations over the distribution $w(\vec{n}_A, \vec{n}_B, \vec{n}_C)$:

$$c_{\alpha\beta\gamma} = \text{tr}(\rho \sigma_\alpha \otimes \sigma_\beta \otimes \sigma_\gamma) = E[(n_A)_\alpha (n_B)_\beta (n_C)_\gamma]. \quad (71)$$

For the ϵ -GHZ state, the coefficients $c_{\alpha\beta\gamma}$ can be read off the Pauli representation in Eq. (69):

$$E[(n_A)_j] = E[(n_B)_j] = E[(n_C)_j] = 0, \quad (72)$$

$$E[(n_A)_j (n_B)_k] = E[(n_A)_j (n_C)_k] = E[(n_B)_j (n_C)_k] = \epsilon \delta_{j3} \delta_{k3}, \quad (73)$$

$$E[(n_A)_j (n_B)_k (n_C)_l] = \epsilon (\delta_{j1} \delta_{k1} \delta_{l1} - \delta_{j1} \delta_{k2} \delta_{l2} - \delta_{j2} \delta_{k1} \delta_{l2} - \delta_{j2} \delta_{k2} \delta_{l1}). \quad (74)$$

Now define a “vector”

$$\vec{N} \equiv ((n_B)_1 (n_C)_1 - (n_B)_2 (n_C)_2) \vec{e}_1 - ((n_B)_1 (n_C)_2 + (n_B)_2 (n_C)_1) \vec{e}_2 + (n_B)_3 \vec{e}_3, \quad (75)$$

whose magnitude satisfies

$$\begin{aligned}\vec{N} \cdot \vec{N} &= ((n_B)_1 (n_C)_1 - (n_B)_2 (n_C)_2)^2 + ((n_B)_1 (n_C)_2 + (n_B)_2 (n_C)_1)^2 + (n_B)_3^2 \\ &= ((n_B)_1^2 + (n_B)_2^2) ((n_C)_1^2 + (n_C)_2^2) + (n_B)_3^2 \\ &\leq \vec{n}_B \cdot \vec{n}_B = 1. \end{aligned} \quad (76)$$

If $w(\vec{n}_A, \vec{n}_B, \vec{n}_C)$ is a probability distribution, we have that

$$5\epsilon = |E[\vec{n}_A \cdot \vec{N}]| \leq E[|\vec{n}_A \cdot \vec{N}|] \leq \frac{1}{2} E[\vec{n}_A \cdot \vec{n}_A + \vec{N} \cdot \vec{N}] \leq 1, \quad (77)$$

which shows that if $\epsilon > 1/5$, the ϵ -GHZ state is nonseparable.

To finish the proof, we need to construct a product ensemble for the ϵ -GHZ state when $\epsilon \leq 1/5$. We begin by defining five density operators

$$\begin{aligned}\rho_1 &\equiv \frac{1}{2} \left(P_{\vec{e}_3} \otimes P_{\vec{e}_3} \otimes P_{\vec{e}_3} + P_{-\vec{e}_3} \otimes P_{-\vec{e}_3} \otimes P_{-\vec{e}_3} \right) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 + 1 \otimes \sigma_3 \otimes \sigma_3 + \sigma_3 \otimes 1 \otimes \sigma_3 + \sigma_3 \otimes \sigma_3 \otimes 1 \right),\end{aligned}\quad (78)$$

$$\begin{aligned}\rho_2 &\equiv \frac{1}{4} \left(P_{\vec{e}_1} \otimes P_{\vec{e}_1} \otimes P_{\vec{e}_1} + P_{\vec{e}_1} \otimes P_{-\vec{e}_1} \otimes P_{-\vec{e}_1} + P_{-\vec{e}_1} \otimes P_{\vec{e}_1} \otimes P_{-\vec{e}_1} + P_{-\vec{e}_1} \otimes P_{-\vec{e}_1} \otimes P_{\vec{e}_1} \right) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 + \sigma_1 \otimes \sigma_1 \otimes \sigma_1 \right),\end{aligned}\quad (79)$$

$$\begin{aligned}\rho_3 &\equiv \frac{1}{4} \left(P_{\vec{e}_1} \otimes P_{\vec{e}_2} \otimes P_{-\vec{e}_2} + P_{\vec{e}_1} \otimes P_{-\vec{e}_2} \otimes P_{\vec{e}_2} + P_{-\vec{e}_1} \otimes P_{\vec{e}_2} \otimes P_{\vec{e}_2} + P_{-\vec{e}_1} \otimes P_{-\vec{e}_2} \otimes P_{-\vec{e}_2} \right) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 - \sigma_1 \otimes \sigma_2 \otimes \sigma_2 \right),\end{aligned}\quad (80)$$

$$\begin{aligned}\rho_4 &\equiv \frac{1}{4} \left(P_{\vec{e}_2} \otimes P_{\vec{e}_1} \otimes P_{-\vec{e}_2} + P_{-\vec{e}_2} \otimes P_{\vec{e}_1} \otimes P_{\vec{e}_2} + P_{\vec{e}_2} \otimes P_{-\vec{e}_1} \otimes P_{\vec{e}_2} + P_{-\vec{e}_2} \otimes P_{-\vec{e}_1} \otimes P_{-\vec{e}_2} \right) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 - \sigma_2 \otimes \sigma_1 \otimes \sigma_2 \right),\end{aligned}\quad (81)$$

$$\begin{aligned}\rho_5 &\equiv \frac{1}{4} \left(P_{\vec{e}_2} \otimes P_{-\vec{e}_2} \otimes P_{\vec{e}_1} + P_{-\vec{e}_2} \otimes P_{\vec{e}_2} \otimes P_{\vec{e}_1} + P_{\vec{e}_2} \otimes P_{\vec{e}_2} \otimes P_{-\vec{e}_1} + P_{-\vec{e}_2} \otimes P_{-\vec{e}_2} \otimes P_{-\vec{e}_1} \right) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 - \sigma_2 \otimes \sigma_2 \otimes \sigma_1 \right),\end{aligned}\quad (82)$$

each of which takes care of one of the nonzero correlations in Eqs. (73) and (74). An equal mixture gives the ($\epsilon = 1/5$)-GHZ state:

$$\begin{aligned}&\frac{1}{5}(\rho_1 + \rho_2 + \rho_3 + \rho_4 + \rho_5) \\ &= \frac{1}{8} \left(1 \otimes 1 \otimes 1 + \frac{1}{5} \left(1 \otimes \sigma_3 \otimes \sigma_3 + \sigma_3 \otimes 1 \otimes \sigma_3 + \sigma_3 \otimes \sigma_3 \otimes 1 \right. \right. \\ &\quad \left. \left. + \sigma_1 \otimes \sigma_1 \otimes \sigma_1 - \sigma_1 \otimes \sigma_2 \otimes \sigma_2 - \sigma_2 \otimes \sigma_1 \otimes \sigma_2 - \sigma_2 \otimes \sigma_2 \otimes \sigma_1 \right) \right),\end{aligned}\quad (83)$$

thus completing the proof.

For both the Werner state and the ϵ -GHZ states, we are able to construct an optimal ensemble based on the 6 cardinal directions, but notice that this ensemble is not the canonical representation of Sec. 2.2.

5 Conclusion

In this paper we introduce a canonical procedure for constructing representations of density operators in terms of complete or overcomplete sets of pure product projectors. The procedure is useful for investigating the separability of density operators: if the expansion coefficients in such a representation are nonnegative, they can be interpreted as a probability distribution, with the result that the representation then provides an explicit product

ensemble for the density operator. Generally, however, the canonical representations have negative values even when the density operator is separable.

The canonical continuous representation is the unique representation that contains only scalar and vector spherical-harmonic terms, and we outline how to obtain all possible pure-product-state representations from the canonical continuous representation by adding arbitrary higher-order-spherical-harmonic content. Searching the space of spherical-harmonic expansions for nonnegative representations would be quite difficult, but we provide an optimal ensemble in the particular case of the ϵ -GHZ state, thereby identifying the separability boundary for this class of states.

Acknowledgments

Thanks to G. J. Milburn and R. Pranaw for discussions of the method used to prove the separability of the ϵ -GHZ state. This work was supported in part by the US Office of Naval Research (Grant No. N00014-93-0-0116) and by the UK Engineering and Physical Sciences Research Council.

Note added

The separability bound for the ϵ -GHZ state derived in Sec. 4 has also been found by Dür, Cirac, and Tarrach [15] as part of a more general result. They prove that the N -qubit ϵ -cat state (51) is separable if and only if $\epsilon < 1/(1 + 2^{N-1})$, which supersedes our Eq. (55).

References

- [1] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
- [2] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 1 (1996).
- [3] K. Życzkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Phys. Rev. A **58**, 883 (1998).
- [4] S. L. Braunstein, C. M. Caves, R. Jozsa, N. Linden, S. Popescu, and R. Schack, submitted to Phys. Rev. Lett. ([quant-ph/9811018](#)).
- [5] D. G. Cory, A. F. Fahmy, and T. F. Havel, Proc. Nat. Acad. Sci. USA **94**, 1634 (1997).
- [6] N. A. Gershenfeld and I. L. Chuang, Science **275**, 350 (1997).
- [7] D. G. Cory, M. D. Price, and T. F. Havel, Physica D **120**, 82 (1998).
- [8] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, Am. J. Phys. **58**, 1131 (1990).
- [9] C. M. Caves, J. Superconductivity, to be published ([quant-ph/9811082](#)).

- [10] C. H. Bennett, D. P. DiVincenzo, C. A. Fuchs, T. Mor, E. Rains, P. W. Shor, J. A. Smolin, and W. K. Wootters, *Phys. Rev. A* **59**, 1070 (1999).
- [11] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).
- [12] M. Murao, M. B. Plenio, S. Popescu, V. Vedral, P. L. Knight, *Phys. Rev. A* **57**, R4075 (1998).
- [13] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, *Phys. Rev. Lett.* **76**, 722 (1996).
- [14] R. Horodecki and M. Horodecki, *Phys. Rev. A* **54**, 1838 (1996).
- [15] W. Dür, J. I. Cirac, and R. Tarrach, [quant-ph/9903018](#).